

Cosmogenic noble gases in silicate inclusions of iron meteorites: Effects of bulk composition on elemental production rates.

O Jentsch & L Schultz

Max-Planck-Institut für Chemie, 55020 Mainz Germany

Abstract

The influence of bulk chemical composition on elemental production rates of neon has been studied experimentally by investigating mineral samples separated from the iron meteorites Four Corners, Landes, Linwood and Zagora. The neon in these samples is purely cosmogenic and its isotopic composition indicates an enhanced production of ^{21}Ne . This is due to the production of additional secondary neutrons caused by a larger multiplicity in meteorites of high-Z bulk chemistry.

Introduction

Meteoroids as meter-sized bodies are irradiated in space by high-energy cosmic-ray particles that penetrate these bodies and interact with its matter. Bauer (1947) and Huntley (1948) independently proposed that measurable amounts of He should be produced in iron meteorites. Paneth *et al.* (1952) showed that this was indeed true and that cosmic-ray produced $^3\text{He}/^4\text{He}$ ratio is in the range 0.2 to 0.3 and were thus very different from atmospheric He. Soon thereafter, also cosmogenic Ne and Ar isotopes were discovered (Reasbeck & Mayne 1955; Gentner & Zähringer 1957). Since then, cosmic-ray produced ("cosmogenic") stable or radioactive reaction products have been studied in great detail to investigate the irradiation history of extraterrestrial matter or the nature of cosmic rays in the past (see *e.g.* Anders 1962; Lal 1972; Reedy *et al.* 1983; Vogt *et al.* 1990).

The production rate of cosmogenic nuclides is dependent on the chemical composition of the target material, the properties of the incident particle irradiation, and the shielding conditions (pre-atmospheric size of the meteoroid and sample's location within the meteoroid). The effects of shielding are mainly caused by the fact that in nuclear reactions initiated by primary cosmic rays secondary particles are also produced. These secondary particles, mostly neutrons of smaller energy, cause further nuclear reactions. Elemental or isotopic nuclide ratios produced by these low-energy particles are different from those of the primary high-energy irradiation. Thus, the shielding of a sample is characterized by elemental or isotopic ratios of specific cosmogenic nuclides produced in different percentages by primary and secondary particles. For example, for shielding corrections of production rates of stony meteorites, cosmogenic $^{22}\text{Ne}/^{21}\text{Ne}$ is commonly used (*e.g.* Eugster 1988), or for iron meteorites, the $^4\text{He}/^{38}\text{Ar}$ ratio (*e.g.* Voshage & Feldmann 1978) is a measure for shielding.

The production rates of cosmogenic nuclides depend twofold on the bulk chemical composition of the meteor-

ite. First, the production of a particular nuclide is given by the concentration-weighted sum of the individual elemental production rates from each target element in the meteorite and, secondly, the elemental production rates themselves depend on it since spectra and intensities of secondary particles inside the meteoroid are influenced by the mean mass number and the mean atomic number of the bulk meteorite. In particular, the multiplicity for the production of secondary neutrons is greater in metal-rich meteorites compared to stony meteorites.

The latter ("matrix") effect was first suggested by Begemann & Schultz (1988). This idea is based on measurements of cosmogenic noble gases in metal and silicates of mesosiderites (Begemann *et al.* 1976). These authors had observed that the concentration of cosmogenic ^{38}Ar produced from Ca in silicates (mainly by secondary neutrons) is larger than expected from the ^{38}Ar produced in metal by the primary irradiation. Begemann and Schultz (1988) show that the production rate ratio of ^{38}Ar from Ca to that of ^{38}Ar from Fe is correlated with the total bulk metal content of mesosiderites. Also, the $^{22}\text{Ne}/^{21}\text{Ne}$ ratio in these silicate samples is relatively low. This is explained by an enhanced production of ^{21}Ne via secondary neutrons and the reaction $^{24}\text{Mg}(n,\alpha)^{21}\text{Ne}$.

The matrix effect was also treated theoretically by Michel *et al.* (1990) and Masarik & Reedy (1994) using model calculations which combine spectra of primary and secondary particles derived from Monte Carlo calculations of intra- and internuclear cascades with thin-target cross sections of the underlying nuclear reactions. These calculations show, for example, that the matrix effect is most pronounced in the production rate of ^{21}Ne from Mg, and compared to stony meteorites the ^{21}Ne production is enhanced by a factor of up to two in silicate inclusions of iron meteorites.

We report here measurements of concentration and isotopic composition of neon in silicates separated from three IA iron meteorites with silicate inclusions (Landes, Linwood and Zagora) and from the IB iron Four Corners. This work was carried out to obtain information on the matrix effect in iron meteorites which are more metal-rich than the previously investigated mesosiderites.

Experimental Methods and Results

Carbon and silicate phases of Landes and Zagora were separated from the meteorite by anodic dissolution of the FeNi. After several ultrasonic treatments and quenching steps with liquid N₂, a further separation of different grain size fractions took place using density and magnetic separation techniques. The Four Corners and Linwood separates were obtained from R S Clarke (Washington DC) and are leftovers from investigations by G P Merrill and E P Henderson.

The concentration of the elements Mg, Al, Si, K, Ca, Ti and Fe of selected separates was determined using x-ray fluorescence (XRF) techniques. The isotopic composition and concentrations of noble gases were measured using apparatus and experimental procedures as described by Schultz *et al.* (1991). Results are given in Table 1 together with estimates of the mineral composition of the samples calculated from its chemical composition.

Figure 1 shows isotopic ratios of neon in a 3-isotope plot. Cosmogenic ²⁰Ne/²²Ne ratios in ordinary chondrites are rather independent of shielding (²⁰Ne/²²Ne = 0.84 ± 0.02) and are found in a narrow area (Fig 1; hatched, marked "Chondrites"). If the measured neon is a mixture of cosmogenic gas and a trapped component (solar or

planetary gas, atmospheric contamination) the data points would be shifted to higher ²⁰Ne/²²Ne values. The low ²⁰Ne/²²Ne values indicate, however, that no significant abundance of trapped Ne is present in spite of trapped Ar, Kr and Xe that is commonly observed in silicate inclusions of iron meteorites (Hintenberger *et al.* 1969; Niemeyer 1979; Crabb 1983; Jentsch & Schultz 1987). The displacement of many data points to higher ²¹Ne/²²Ne values is discussed below.

The three feldspar-rich samples are characterized by smaller ²⁰Ne/²²Ne and ²¹Ne/²²Ne ratios. This is due to the relatively high concentration of Na in these samples. Sodium is an element that has a high production rate for ²²Ne compared to that of the other two neon isotopes. Smith & Huneke (1975) observed ²⁰Ne/²²Ne ratios < 0.7 in plagioclase separated from ordinary chondrites. Our feldspar measurement of Zagora is very similar to this value. The two Landes samples with a smaller percentage of feldspar (and thus Na) are located on the mixing line between the "pure" feldspar neon ratio and "normal" cosmogenic neon. Unfortunately, the concentration of Na was not measured directly in these samples because NaNO₃ was used to make the tablets for XRF-measurements.

Table 1.

Chemical and mineralogical information on the investigated separates as well as on their concentration and isotopic composition of neon.

Meteorite and sample #	Mg	Al	Si	K	Ca	Ti	Fe	Mg/(Mg+Al+Si) [wt.% ratio]	Mineral composition				²⁰ Ne	²¹ Ne	²² Ne	²² Ne/ ²¹ Ne [in 10 ⁻³ cm ³ STP/g]
	[wt %]								Opx	Ol	Cpx	Fsp	[wt.%]			
Four Corners																
401	19.5	1.22	26.5	0.075	2.50	0.13	3.8	0.414	67	12	12	9	348.1	407.9	394.6	0.967
Landes																
I364	22.4	0.11	26.0	0.022	1.20	0.12	4.0	0.462	81	14	5	-	78.5	89.4	91.3	1.021
I344	21.9	0.35	25.7	0.021	1.59	0.13	4.3	0.456	79	14	7	-	75.2	85.5	88.0	1.029
I363	26.9	0.09	23.0	0.016	.78	0.07	3.8	0.539	42	54	4	-	84.8	97.7	98.5	1.008
I355	4.6	9.90	31.2	0.36	1.95	0.07	2.9	0.100	22	-	-	78	15.2	17.1	19.9	1.161
II345	22.2	0.37	25.6	0.027	1.42	0.13	4.3	0.460	78	15	7	-	79.7	91.2	92.8	1.017
II365	23.1	0.23	25.3	0.022	1.14	0.17	4.1	0.474	74	21	5	-	60.5	68.9	70.3	1.021
II366	26.0	0.11	23.7	0.015	0.69	0.08	3.7	0.522	53	44	3	-	89.9	103.2	103.8	1.006
II353	5.4	9.40	31.2	0.33	1.98	0.07	2.6	0.117	26	-	-	74	20.2	22.7	26.4	1.162
III343	21.4	0.44	25.1	0.021	2.38	0.13	5.0	0.456	70	17	13	-	75.3	85.8	87.6	1.021
III347	25.2	0.21	23.7	0.010	1.49	0.10	3.9	0.513	49	43	8	-	86.5	99.8	100.9	1.011
Linwood																
425	11.5	3.49	30.8	0.21	3.23	0.14	2.7	0.252	43	-	13	44	32.3	36.6	36.1	0.987
Zagora																
I329	23.0	0.22	25.4	0.015	1.12	0.13	4.1	0.473	76	19	5	-	91.6	108.2	104.3	0.964
I417	24.4	0.15	24.6	0.015	0.99	0.12	3.9	0.496	65	31	4	-	118.0	139.1	133.5	0.960
I328	22.2	0.24	25.5	0.023	1.21	0.14	4.8	0.464	63	32	5	-	113.7	133.7	129.2	0.966
II322	23.5	0.07	24.3	0.013	0.81	0.11	5.9	0.490	70	27	3	-	110.8	130.2	125.4	0.963
II320	24.3	0.09	24.1	0.017	0.86	0.11	5.1	0.501	64	33	3	-	110.1	129.9	124.0	0.955
II319	23.7	-	23.3	0.026	1.05	0.10	7.1	0.504	59	36	5	-	112.6	132.6	127.5	0.962
II325	25.3	0.08	22.8	0.020	2.54	0.09	4.3	0.525	33	52	15	-	119.6	141.9	135.1	0.952
III386	0.94	11.8	32.8	0.63	1.82	0.05	2.1	0.021	9	-	-	91	20.8	23.1	35.2	1.523

Figure 4 (adopted from Begemann & Schultz 1988) shows $^{22}\text{Ne}/^{21}\text{Ne}$ as a function of the $\text{Mg}/(\text{Mg}+\text{Si}+\text{Al})$ ratio. More than 95% of cosmogenic neon is produced in chondrites under normal shielding conditions from these three elements, about 80% alone from Mg. There is a small dependence of the $^{22}\text{Ne}/^{21}\text{Ne}$ on the chemical composition as indicated by the lines connecting measurements of mineral separates of individual meteorites. Referring to the same chemistry [= $\text{Mg}/(\text{Mg}+\text{Si}+\text{Al})$] the lowest $^{22}\text{Ne}/^{21}\text{Ne}$

Figure 4. Cosmogenic $^{22}\text{Ne}/^{21}\text{Ne}$ ratios of silicates from chondrites and mesosiderites that have values smaller than 1.06 (adapted from Begemann & Schultz 1988). Data points are connected when different mineral separates from the same meteorite have been analyzed.

Figure 5. $^{22}\text{Ne}/^{21}\text{Ne}$ ratios of silicate separates from iron meteorites as function of their $\text{Mg}/(\text{Mg}+\text{Si}+\text{Al})$ ratio. The position of chondrites is shown by the hatched area. For the same chemical composition the silicates of iron meteorites have lower $^{22}\text{Ne}/^{21}\text{Ne}$ ratios due to an enhanced flux of secondary neutrons. Samples with low Mg concentrations are feldspar-rich with relatively high sodium contents. Their $^{22}\text{Ne}/^{21}\text{Ne}$ is influenced by higher production rates of ^{22}Ne from Na.

It should also be noted that in the metal phase of iron meteorites, cosmogenic $^{22}\text{Ne}/^{21}\text{Ne}$ is generally greater than 1.06 (see compilation by Schultz & Kruse 1989).

References

- Anders E 1962 Meteorite Ages. *Reviews of Modern Physics* 34:287-325.
- Bauer C A 1947 Production of helium in meteorites by cosmic radiation. *Physical Review* 72:354-355.
- Begemann F & Schultz L 1988 The influence of bulk chemical composition on the production rate of cosmogenic nuclides in meteorites. *Lunar and Planetary Science* 19:51-52.
- Begemann F, Weber H W, Vilcsek E & Hintenberger H 1976 Rare gases and ^{36}Cl in stony-iron meteorites: Cosmogenic elemental production rates, exposure ages, diffusion losses and thermal histories. *Geochimica et Cosmochimica Acta* 40:353-368.
- Begemann F, Zhaohui Li, Schmitt-Strecker S, Weber H W & Zitu Xu 1985 Noble gases and the history of Jilin meteorite. *Earth and Planetary Science Letters* 72:247-262.
- Begemann F, Fan Caiyun, Weber H W & Wang Xianbin 1996 Light noble gases in Jilin: More of the same and something new. *Meteoritics and Planetary Science*: *in press*.
- Crabb J 1983 On the siting of noble gases in silicate inclusions of the El Taco iron Meteorite. *Lunar and Planetary Science* 14:123-135.
- Eberhardt P, Eugster O, Geiss J & Marti K 1966 Rare gas measurements in 30 stone meteorites. *Zeitschrift für Naturforschung* 21a:416-426.
- Eugster O 1988 Cosmic-ray production rates for ^3He , ^{21}Ne , ^{38}Ar , ^{83}Kr , and ^{126}Xe in chondrites based on ^{81}Kr -Kr ages. *Geochimica et Cosmochimica Acta* 52:1649-1662.
- Gentner W & Zähringer J 1957 Argon und Helium als Kernreaktionsprodukte in Meteoriten. *Geochimica et Cosmochimica Acta* 11:60-71.
- Hintenberger H, Schultz L & Weber H 1969 Rare gases in the iron and in the inclusions of the Campo del Cielo meteorite, El Taco. In: *Meteorite Research* (ed P M Millman). Reidel, Dordrecht, 895-900.
- Huntley H E 1948 Production of helium by cosmic rays. *Nature* 161:356.
- Jentsch O & Schultz L 1987 Trapped noble gases in silicate inclusions of the Landes iron meteorite. *Meteoritics* 22:418-419.
- Lal D 1972 Hard rock cosmic archeology. *Space Science Reviews* 14:3-102.
- Loeken T, Scherer P, Weber H W & Schultz L 1992 Noble gases in eighteen stone meteorites. *Chemie der Erde* 52:249-259.
- Masarik J & Reedy R C 1994 Effects of bulk composition on nuclide production processes in meteorites. *Geochimica et Cosmochimica Acta* 58:5307-5317.
- Michel R, Dragovitsch P & Filges D 1990 On the dependence of cosmogenic nuclide production rates in meteoroids on bulk chemical composition. *Meteoritics* 25:386-387.
- Nagai H, Honda M, Imamura M & Kobayashi K 1993 Cosmogenic ^{10}Be and ^{26}Al in metal, carbon, and silicate of meteorites. *Geochimica et Cosmochimica Acta* 57:3705-3723.
- Niemeyer S 1979 I-Xe dating of silicate and troilite from IAB iron meteorites. *Geochimica et Cosmochimica Acta* 43:843-860.
- Paneth F A, Reasbeck P & Mayne K I 1952 Helium 3 content and age of meteorites. *Geochimica et Cosmochimica Acta* 2:300-303.
- Reasbeck P & Mayne K I 1955 Cosmic radiation effects in meteorites. *Nature* 176:733-734.
- Reedy R C, Arnold J R & Lal D 1983 Cosmic-ray record in solar system matter. *Annual Review of Nuclear Particle Science* 33:505-557.
- Schultz L & Kruse H 1989 Helium, neon, and argon in meteorites - a data compilation. *Meteoritics* 24:155-172.
- Schultz L, Weber H W & Begemann F 1991 Noble gases in H-chondrites and potential differences between Antarctic and non-Antarctic meteorites. *Geochimica et Cosmochimica Acta* 55:59-66.
- Smith S P & Huneke J C 1975 Cosmogenic neon produced from sodium in meteoritic minerals. *Earth and Planetary Science Letters* 27:191-199.
- Vogt S, Herzog G F & Reedy R C 1990 Cosmogenic nuclides in extraterrestrial materials. *Reviews of Geophysics* 28:253-275.
- Voshage H & Feldmann H 1978 Investigations on cosmic-ray-produced nuclides in iron meteorites. 1. The measurement and interpretation of rare gas concentrations. *Earth and Planetary Science Letters* 39:25-36.