

Aquatic fauna of the Warren bioregion, south-west Western Australia: Does reservation guarantee preservation?

K M Trayler¹, J A Davis¹, P Horwitz² & D Morgan¹

¹School of Biological and Environmental Sciences, Murdoch University, Murdoch WA 6150;

²Centre for Ecosystem Management, Edith Cowan University, Joondalup WA 6027

Abstract

The Warren Bioregion, in the extreme south-west of Western Australia, has a unique assemblage of aquatic invertebrates, fish and amphibians. Current literature indicates that 192 fully described species have been collected, of which 10 invertebrate, 1 fish and 6 frog species could be considered locally endemic. We estimate that secure nature reserves (A-Class and National Parks) in the Warren Bioregion provide a refuge for 86% of the aquatic faunal elements. Reservation alone, however, may not be sufficient to protect certain of the aquatic fauna. Adverse impacts occurring within catchments, including erosion and deposition of sediment, salinization, fire, land clearing, the presence of dams and the introduction of exotic fish, may adversely affect the aquatic fauna within a reserve. Management of protected habitats must ensure that only anthropogenic activities which are sympathetic to the long term persistence of all elements of the biota occur within, and adjacent to, the reserve systems.

Introduction

The natural landscape of the south-western corner of Australia is characterised by rivers which arise on an ancient and flat semi-arid inland plateau. From here, the rivers flow sluggishly towards the coast, before passing through a zone where the topography steepens and rainfall increases (Mulcahy *et al.* 1972; Churchward *et al.* 1988). Beyond this, the rivers again slow as they traverse the coastal lowlands, which feature extensive wetland systems and terminate in lagoon-like estuaries. Early settlers found the coastal lowlands to be largely infertile soils and used much of the area as pastoral land, preferring to use the rich alluvial soils along the rivers for intensive agriculture (Jarvis 1979). Pressures of urbanization and intensification of agriculture, associated with a burgeoning population in the century since colonization, have led to the loss of a large proportion of coastal wetland systems and many of those that remain have been altered from their natural state (see Halse 1989). A number of the larger rivers have been damaged through siltation and salinization, and many of the remainder are now impounded or subject to varied degrees of alteration through activities such as mining and logging (Olsen & Skitmore 1991).

The effect of these changes on the aquatic fauna cannot be fully assessed due to the lack of historical or baseline data. Locally or even regionally endemic aquatic species may have been lost from south-western Australia, or suffered substantial range reductions, particularly where these fauna were sensitive to changed hydrological regimes, increased eutrophication and/or salinization. This is evident in our aquatic environments today. Highly eutrophic wetlands contain few rare taxa,

have high densities of "pollution tolerant" species and, overall, support an invertebrate composition which is distinct from that occurring in low to moderately enriched or coloured wetlands (Davis *et al.* 1993; Edward *et al.* 1994). The highly saline Hotham River and Thirty-Four Mile Brook, south-east of Perth, are dominated by salt tolerant crustaceans (60% total abundance), while insects contribute only 14% to the overall invertebrate composition (Bunn & Davies 1992). This situation is atypical of undisturbed streams of the jarrah forest, where insects usually comprise 70-80% of the fauna (Bunn *et al.* 1986), and was attributed to increased salinity (Bunn & Davies 1992). Williams *et al.* (1991) found little or no longitudinality in the faunal composition of the Blackwood River despite the presence of a distinct salinity gradient along its length. Although this could be interpreted as evidence of a halotolerant fauna, it may also indicate elimination of less tolerant fauna, which once characterised a more diverse system.

Public awareness of the need to conserve aquatic habitats has increased markedly over the last decade and this is reflected in the number of scientific studies that have been undertaken on wetlands, streams and rivers in the south-west since the mid 1980s (see Balla 1994). Considerable attention has been devoted to the aquatic fauna and their habitats in highly populated areas of the south-west, such as wetlands on the Swan Coastal Plain and streams in the adjacent jarrah forest, where monitoring to assess the impact of pollution or habitat change has often been the principal research or management objective (*e.g.* Storey *et al.* 1990; Bunn & Davies 1992; Growns *et al.* 1992).

Further south, the relatively undisturbed aquatic habitats within the Warren Bioregion (*sensu* Thackway & Cresswell 1995) have also provided a focus for scientific studies (*e.g.* Christensen 1982; Pusey & Edward 1990a, b; Horwitz 1996; Edward *et al.* 1994). This region is regarded as an important centre for endemism from a

botanical viewpoint (Hopper *et al.* 1992), but its significance to the aquatic fauna has not been assessed in its entirety. This paper summarizes existing knowledge of the aquatic fauna in the Warren Bioregion and estimates the degree of endemism found in the region. In addition, it presents a preliminary assessment of the reservation status of known aquatic species and makes recommendations for the ongoing protection of fauna within the reserve system.

Approaches

The aquatic invertebrates (>110 µm), fishes and amphibians which occur within the Warren Bioregion were compiled from published literature and museum records, and are therefore limited by the habitats examined and sampling protocol of these sources. The distribution of each species, determined from published literature, is described as either widespread, regionally endemic or locally endemic in accordance with Horwitz (1996). The reservation status of each species within the Warren Bioregion has been determined according to their presence or absence within reserves which fall under categories I and II of the World Conservation Union's classification of protected areas (Anon. 1994). This equates only to nature reserves (A-Class) and national parks which have the conservation of flora and fauna as a primary objective and require legislative provisions to alter their boundaries or status. Other forms of reserves in the region do not have this level of security and therefore have not been included in this study. We do, however, acknowledge that these other areas also play an important role in conservation. For the purpose of this study, a species which is described as reserved must have been recorded at least once in a secure reserve, as defined above.

To date, there have been no comprehensive surveys of aquatic invertebrates within all nature reserves and national parks of the Warren Bioregion and therefore the reference in Table 1 to species as absent from these reserve categories should be considered preliminary. In addition, species which are widespread or regionally endemic may not be found in nature reserves or national parks within this region, but may be found within similar reserves located elsewhere.

Although more invertebrate taxa are found in the Warren Bioregion than have been recorded in Table 1, many of these have not yet been described and specimens only exist in voucher or reference collections of individual researchers. To prevent confusion, whereby authors may have assigned different voucher names to the same taxon, only species for which published descriptions exist were considered here. Emergent adult invertebrates found in the vicinity of aquatic habitats were excluded because they could not be definitively linked with specific waterbodies. In addition, waterbirds have been excluded from this review, but information relating to their occurrence and reservation within the larger southern forest region may be found in Christensen (1992).

The tabulated information on aquatic invertebrates, fishes and amphibians provides a preliminary account of the level of endemism of aquatic fauna in the Warren

Bioregion and emphasises the importance of the reserve system to both the restricted and more common fauna of the region.

Local endemism in the aquatic fauna of the Warren Bioregion

Invertebrates

Aquatic habitats within the Warren Bioregion support a large proportion of invertebrate taxa which are endemic to south-west Western Australia, with approximately 17% of these considered locally restricted on the basis of surveys (Table 1). This number, however, may be an underestimate as only fully-described species have been considered, thereby excluding some taxa. There are, for example, only eight oligochaete species with widespread distributions shown in Table 1, yet there are at least a further four species which are restricted to the Warren Bioregion (Horwitz 1996) for which descriptions are pending (A Pinder *pers. comm.*). This situation may also extend to other groups, notably the dipterans and arachnids. Locally endemic invertebrates include the freshwater crayfish *Engaewa subcoerulea* and *E. similis* and the trichopteran *Kosrheithrus boorarus* (see Horwitz 1994; Grouns & Davis 1994, respectively).

A rich suite of micro-crustaceans have been found in the area and, in the case of the copepods, this is highly distinctive (Table 1). Bayly (1992) examined temporary ponds near Northcliffe and identified several copepods which are locally endemic, including *Calamoecia elongata*, *Boeckella geniculata* and *Paracyclops* sp nov, as well as a unique form of *C. tasmanica* sl. In addition, the copepod *Hemiboeckella powellensis* is known only from Lake Powell, between Albany and Denmark (Bayly, 1979). Locally endemic cladocerans, *Biaptera imitaria* and *Daphnia occidentalis* were also found near Northcliffe by Bayly (1992). *Daphnia occidentalis* is a relictual species thought to have originated 70 mybp (Benzie 1986, 1988). This species was found in large numbers in a single pool, surrounded by swamp land within the D'Entrecasteaux National Park.

Fish

Eight species of freshwater fish are endemic to the south-west of Western Australia (Table 2) and four of these are now principally confined to the Warren Bioregion (Morgan *et al.* 1996). These include the Western Australian salamanderfish *Lepidogalaxias salamandroides*, a relictual Gondwanaland species (Allen 1982), which is now predominantly restricted to ephemeral pools within the coastal peat flats between Windy Harbour and Walpole (Morgan *et al.* 1996). The black stripe minnow *Galaxiella nigrostriata* and Balston's pygmy perch *Nannatherina balstoni* are similarly confined, although the former species is found in low numbers in a few lakes of the region, while the latter occurs in low numbers in both lakes and rivers of the region (Morgan *et al.* 1996). The mud minnow *Galaxiella munda* has the widest distribution of these four species, occurring in the headwaters and tributaries of rivers, as well as on the coastal peat flats (Morgan *et al.* 1996).

Disjunct and isolated populations of some of these

Table 1

Invertebrate fauna found in aquatic habitats within the Warren bioregion. Reservation status refers to their presence or absence in a nature reserve or national park within the bioregion. Distribution: W, widespread; RE, regionally endemic; LE, locally endemic. Sources: 1, Pusey & Edward (1990a); 2, Grouns & Davis (1994); 3, Edward *et al.* (1994); 4, Bayly (1982); 5, Horwitz (1994); 6, Williams *et al.* (1991); 7, Bayly (1992); 8, Bayly (1979); 9, Harvey (1996); 10, Morton (1990).

Species ¹	Source	Lotic	Lentic	Reservation	Distribution
MOLLUSCA					
Hydriidae					
<i>Westralunio carteri</i> Iredale	2,3	√	√	+	RE
Glacidorbidae					
<i>Glacidorbis occidentalis</i> Bunn & Stoddart	1	√		+	RE
Ancylidae					
<i>Ferrissia petterdi</i> Johnston	1		√	+	W
Hydrobiidae					
<i>Potamopyrgus niger</i> Quoy & Gaimard	6		√	-	W
ANNELIDA					
Phreodrilidae					
<i>Insulodrilus lacustris</i> Benham	5		√	+	W
<i>Insulodrilus nudus</i> Brinkhurst & Fulton	5		√	+	W
Tubificidae					
<i>Limnodrilus hoffmeisteri</i> Cleperede	5		√	+	W
Naididae					
<i>Pristina longiseta</i> Ehrenberg	5		√	+	W
<i>Pristina aequiseta</i> Bourne	5		√	-	W
<i>Dero furcatus</i> Muller	5		√	-	W
<i>Dero digitata</i> Muller	5		√	+	W
<i>Chaetogaster diastrophus</i> Gruithuisen	5		√	-	W
ARACHNIDA					
Hydryphantidae					
<i>Pseudohydryphantes doegi</i> Harvey	5		√	+	RE
Aturidae					
<i>Wheenyoides cooki</i> Harvey	5		√	+	RE
Limnocharidae					
<i>Limnocharis australica</i> Lundblad	5		√	+	W
Pionidae					
<i>Australotiphys barmutai</i> Harvey	9		√	+	RE
<i>Larri laffa</i> Harvey	9		√	+	RE
<i>Piona cumberlandensis</i> Rainbow	9		√	+	W
Arrenuridae					
<i>Arrenurus</i> sp nr <i>tasmanicus</i> Lundblad	5		√	+	RE
OSTRACODA					
Limnocytheridae					
<i>Limnocythere mowbrayensis</i> Chapman	1,3	√	√	+	W
<i>Gomphodella</i> aff <i>maia</i> De Deckker	3		√	+	W
Cyprididae					
<i>Candonocypris novaezelandiae</i> Baird	2,3	√	√	+	W
<i>Newnhamia fenestra</i> King	3		√	+	W
<i>Cypretta viridis</i> Thomson	7		√	+	W
<i>Cypretta baylyi</i> McKenzie	3,4,7		√	+	W
<i>Eucypris virens</i> Jurine	7		√	+	W
<i>Kapcypridopsis asymmetra</i> De Deckker	4		√	+	LE
<i>Sarcypridopsis aculeata</i> Costa	3		√	+	W
<i>Alboa wooroa</i> De Deckker	3		√	+	W
<i>Ilyodromus condonites</i> De Decker	4		√	+	LE
<i>Ilyodromus varrovillus</i> King	7		√	+	W
<i>Bennelongia australis</i> Brady	8		√	+	W
Candonidae					
<i>Candonopsis tenuis</i> Brady	3		√	+	RE
COPEPODA					
Centropagidae					
<i>Calamoecia attenuata</i> Fairbridge	1,3,7	√	√	+	RE
<i>Calamoecia tasmanica</i> Smith sl	1,3,7	√	√	+	W
<i>Calamoecia elongata</i> Bayly	7		√	+	LE
<i>Hemiboeckella searli</i> Sars	3,7		√	+	W
<i>Hemiboeckella andersonae</i> Bayly	3,7		√	+	RE
<i>Hemiboeckella powellensis</i> Bayly	8		√	+	LE
<i>Boeckella geniculata</i> Bayly	7		√	+	LE

Species	Source	Lotic	Lentic	Reservation	Distribution
<i>Gladioferens imparipes</i> Thomson	3		√	+	RE
Cyclopidae					
<i>Macrocyclops albidus</i> Jurine	1,7	√	√	+	W
<i>Australocyclops australis</i> Sars	7		√	-	W
<i>Eucyclops spatulatus</i> Morton	10		√	-	W
CLADOCERA					
Chydoridae					
<i>Biapertura</i> cf. <i>macrocopa</i> Sars	4,7		√	+	W
<i>Biapertura rigidicaudis</i> Smirnov	4		√	+	W
<i>Biapertura affinis</i> Leydig	1	√	√	+	W
<i>Biapertura longinqua</i> Smirnov	7		√	+	W
<i>Biapertura</i> nr. <i>setigera</i> Brehm	3		√	+	W
<i>Biapertura imitatoria</i> Smirnov	7		√	+	LE
<i>Graptoleberis testudinaria</i> Fischer	3		√	+	W
<i>Camptocercus</i> cf. <i>australis</i> Sars	3		√	+	W
<i>Chydorus barroisi</i> Richard	4		√	+	W
<i>Chydorus</i> cf. <i>sphaericus</i> O F Muller	7		√	-	W
<i>Pleuroxus inermis</i> Sars	7		√	-	W
<i>Pleuroxus jugosus</i> Henry	4		√	+	W
<i>Alonella</i> cf. <i>excisa</i> Fischer	7		√	+	W
<i>Rak obustus</i> Smirnov & Timms	8		√	-	W
<i>Monope reticula</i> Henry	7		√	+	RE
Macrothricidae					
<i>Neothrix armata</i> Gurney	1,3,4,7	√	√	+	W
Daphniidae					
<i>Simocephalus acutirostratus</i> King	1,3	√	√	+	W
<i>Daphnia carinata</i> King sl	3		√	+	W
<i>Daphnia occidentalis</i> Benzie	7		√	+	LE
<i>Scapholeberis kingi</i> Sars	7		√	+	W
Bosminidae					
<i>Bosmina meridionalis</i> Sars	3		√	+	W
ISOPODA					
Amphisopidae					
<i>Amphisopus annectans</i> Nicholls	1,5		√	+	RE
<i>Amphisopus ?lintoni</i> Nicholls	1	√	√	+	RE
<i>Hyperodesipus ?plumosus</i> Nicholls & Milner	2	√		-	RE
AMPHIPODA					
Perthidae					
<i>Perthia acutitelson</i> Straskraba	1,2	√	√	+	RE
<i>Perthia branchialis</i> Nicholls	3		√	-	RE
Ceinidae					
<i>Austrochiltonia subtenuis</i> Sayce	3		√	+	W
DECAPODA					
Parastacidae					
<i>Cherax quinquecarinatus</i> Gray	3,5		√	+	RE
<i>Cherax tenuimanus</i> Smith	3,5		√	+	RE
<i>Cherax preissii</i> Erichson	5		√	+	RE
<i>Cherax crassimanus</i> Riek	5		√	+	RE
<i>Cherax destructor</i> Clark	3		√	+	W
<i>Engaewa subcoerulea</i> Riek	5		√	+	LE
<i>Engaewa similis</i> Riek	5		√	+	LE
Grapsidae					
<i>Leptograpsodes octodentatus</i> Milne-Edwards	5		√	+	W
Palaemonidae					
" <i>Palaemonetes</i> " <i>australis</i> Dakin	3,5		√	+	RE
ANISOPTERA					
Aeshnidae					
<i>Aeshna brevistyla</i> Rambur	3		√	+	W
<i>Austroaeschna anacantha</i> Tillyard	2,5,6	√	√	+	RE
Corduliidae					
<i>Orthetrum caledonicum</i> Brauer	3,5,6	√	√	+	W
<i>Austrothemis nigrescens</i> Martin	3,5		√	+	W
<i>Diplacodes bipunctata</i> Brauer	5		√	-	W
<i>Nannophya dalei occidentalis</i> Tillyard	5		√	+	RE
<i>Lathrocordulia metallica</i> Tillyard	3		√	+	RE
<i>Hesperocordulia berthoudi</i> Tillyard	3		√	+	RE

Species	Source	Lotic	Lentic	Reservation	Distribution
<i>Hemicordulia australiae</i> Rambur	3,6	√	√	+	W
<i>Hemicordulia tau</i> Selys	5		√	+	W
<i>Procordulia affinis</i> Selys	3,5		√	+	RE
<i>Synthemis cyanitincta</i> Tillyard	2,3,5	√	√	+	RE
Gomphidae					
<i>Hemigomphus armiger</i> Tillyard	2	√		-	RE
<i>Austrogomphus lateralis</i> Selys	2,3	√	√	+	RE
<i>Austrogomphus collaris</i> Hagen	3,5		√	+	W
<i>Austrogomphus ochraceus</i> Selys	6	√		-	W
ZYGOPTERA					
Lestidae					
<i>Austrolestes annulosus</i> Selys	3,5		√	+	W
<i>Austrolestes analis</i> Rambur	5		√	+	W
Megapodagriidae					
<i>Argiolestes minimus</i> Tillyard	5		√	+	RE
<i>Austroagrion cyane</i> Selys	3		√	+	W
<i>Xanthagrion erythronurum</i> Selys	6	√		-	W
EPHEMEROPTERA					
Leptophlebiidae					
<i>Bibulmena kadjina</i> Dean	1,2,3	√	√	+	RE
<i>Neboissophlebia occidentalis</i> Dean	3		√	+	W
<i>Nyungara bunni</i> Dean	1,2	√		+	RE
<i>Nyungara ellitasha</i> Dean	2	√		-	W
Caenidae					
<i>Tasmanocoenis tillyardi</i> Lestage	2,3	√	√	+	W
PLECOPTERA					
Gripopterygidae					
<i>Newmanoperla exigua</i> Kimmins	1,2	√	√	+	RE
<i>Leptoperla australica</i> Enderlain	1,2	√		+	RE
MEGALOPTERA					
Corydalidae					
<i>Archichauliodes cervulus</i> Theischinger	2	√		-	W
TRICHOPTERA					
Leptoceridae					
<i>Lectrides parilis</i> Neboiss	1,2	√	√	+	RE
<i>Triplectides australis</i> Navas	3		√	+	W
<i>Condocerus nr aptus</i> Neboiss	2	√		-	RE
<i>Notoperata tenax</i> Neboiss	3		√	+	RE
Atriplectididae					
<i>Atriplectides dubius</i> Mosely	2,3	√	√	+	W
Hydroptilidae					
<i>Acroptila globosa</i> Wells	1,3		√	+	RE
Ecnomidae					
<i>Ecnomina ?trulla</i> Neboiss	1,3	√	√	+	RE
<i>Ecnomus pansus</i> Neboiss	1,3		√	+	W
<i>Ecnomus turgidus</i> Neboiss	3		√	-	W
Philorheithridae					
<i>Kosrheithrus boorarus</i> Neboiss	2	√		-	LE
Hydrobiosidae					
<i>Apsilochorema urdalum</i> Neboiss	2	√		-	RE
<i>Taschorema pallescens</i> Banks	2	√		-	RE
Hydropsychidae					
<i>Smicrophylax australis</i> Ulmer	2	√		-	RE
Polycentropodidae					
<i>Plectrocnemia eximia</i> Neboiss	3		√	+	W
<i>Adectophylax volutus</i> Neboiss	2	√		-	RE
COLEOPTERA					
Dytiscidae					
<i>Sternopriscus ?browni</i> Sharp	1,3	√	√	+	RE
<i>Sternopriscus marginatus</i> Watt	1	√		+	RE
<i>Homoeodytes scutellaris</i> Germar	1,3		√	+	W
<i>Rhantus suturalis</i> MacLeay	1	√	√	+	W
<i>Liodessus inornatus</i> Sharp	3		√	+	RE
<i>Liodessus dispar</i> Sharp	3		√	+	RE

Species	Source	Lotic	Lentic	Reservation	Distribution
<i>Megaporus solidus</i> Sharp	3		√	+	RE
<i>Necterosoma darwini</i> Babington	3		√	+	RE
<i>Antiporus femoralis</i> Boheman	3		√	+	W
<i>Lancetes lanceolatus</i> Clark	3		√	+	W
DIPTERA					
Simuliidae					
<i>Austrosimulium furiosum</i> Skuse	1,5	√	√	+	W
<i>Cnephia tonnoiri tonnoiri</i> Drummond	5		√	-	W
Chironomidae					
<i>Aphroteniella filicornis</i> Brundin	1,2,3,5	√	√	+	W
<i>Aphroteniella tenuicornis</i> Brundin	5		√	-	W
<i>Paramerina levidensis</i> Skuse	1,2,3,5	√	√	+	W
<i>Procladius palludicola</i> Skuse	3,5		√	+	W
<i>Procladius ?villosimanus</i> Kieffer	3		√	+	W
<i>Alotanypus dalyupensis</i> Freeman	1,3,5		√	+	W
<i>Coelopynia pruinosa</i> Freeman	3		√	+	W
<i>Corynoneura ?scutellata</i> Winnertz	5		√	+	W
<i>Stictocladus uniserialis</i> Freeman	1,5	√	√	+	W
<i>Cricotopus annuliventrus</i> Skuse	1,2,3,5	√	√	+	W
<i>Paralimnophyes pullulus</i> Skuse	3,5		√	+	W
<i>Cladopelma curtivalva</i> Kieffer	1,3,5	√	√	+	W
<i>Cryptochironomus griseidorsum</i> Kieffer	1,3,5	√	√	+	W
<i>Chironomus occidentalis</i> Skuse	3		√	+	W
<i>Chironomus aff alternans</i> Walker	1,3,5		√	+	W
<i>Chironomus tepperi</i> Skuse	4		√	+	W
<i>Dicrotendipes ?conjunctus</i> Walker	1,3,5	√	√	+	W
<i>Kiefferulus martini</i> Freeman	1,3,5	√	√	+	W
<i>Kiefferulus intertinctus</i> Skuse	3,5		√	+	W
<i>Paratanytarsus grimmii</i> Schneider	5		√	+	W
<i>Stempellina ?australiensis</i> Freeman	3,5		√	+	W

¹ fully described species only

Table 2

Native fish species found in freshwater habitats in the Warren Bioregion. Reservation refers to their presence or absence within a nature reserve (A-class) or national park within the region. Distribution: W, widespread; RE, regionally endemic; LE, locally endemic. Sources of information include Christensen (1982), Jaensch (1992), Morgan *et al.* (1996) and Western Australian Museum records.

Species	Lotic	Lentic	Reservation	Distribution
AGNATHA				
Geotriidae				
<i>Geotria australis</i> Gray	√		+	W
TELEOSTEI				
Lepidogalaxiidae				
<i>Lepidogalaxias salamandroides</i> Mees		√	+	LE
Galaxiidae				
<i>Galaxiella nigrostriata</i> Shipway		√	+	RE
<i>Galaxiella munda</i> McDowall	√	√	+	RE
<i>Galaxias occidentalis</i> Ogilby	√	√	+	RE
<i>Galaxias truttaceus</i> Cuvier	√	√	+	W
<i>Galaxias maculatus</i> Jeyns	√	√	+	W
Percichthyidae				
<i>Bostockia porosa</i> Castelnau	√	√	+	RE
Nannopercidae				
<i>Edelia vittata</i> Castelnau	√	√	+	RE
<i>Nannatherina balstoni</i> Regan	√	√	+	RE
Plotosidae				
<i>Tandanus bostocki</i> Whitley	√	√	+	RE
Gobiidae				
<i>Pseudogobius olorum</i> Sauvage	√	√	+	W ¹
<i>Afurcagobius suppositus</i> Sauvage	√	√	+	RE ¹
Atheriniidae				
<i>Leptatherina wallacei</i> Prince, Ivantsoff & Potter	√	√	+	RE ¹

¹ not strictly a freshwater species.

species have been recorded from Margaret River, Bunbury, Gingin and Two Peoples Bay (Museum Records; Morgan *et al.* 1996) suggesting that they were once more widely distributed throughout the coastal environment of south-west Western Australia. All of these fishes are either included or have been recommended for inclusion in the list of Australian threatened fishes (Anon. 1994).

Amphibians

Twenty two species of frogs are found in the Warren Bioregion, with three of these occurring just within its boundaries (Table 3). Six of the remaining eighteen species could be considered to be locally endemic, including the south coast froglet *Ranidella subinsignifera*, the roseate frog *Geocrinia rosea*, the Walpole frog *G. lutea*, the white-bellied frog *G. alba*, the orange-bellied frog *G.*

Table 3

Frog species known to occur within the boundaries of the Warren Bioregion. Reservation refers to their presence or absence in a nature reserve (A-class) or national park within the region. Distribution: RE, regionally endemic; LE, locally endemic. Sources of information include Main (1965), Christensen (1992), Tyler *et al.* (1994), Roberts *et al.* (in press) and Wardell-Johnson (*pers. comm.*).

Species	Reservation	Distribution
Hylidae		
<i>Litoria adelaidensis</i> Gray	+	RE
<i>Litoria moorei</i> Copeland	+	RE
Myobatrachidae		
<i>Limnodynastes dorsalis</i> Gray	+	RE
<i>Heleioporus inornatus</i> Lee & Main	+	RE
<i>Heleioporus eyrei</i> Gray	+	RE
<i>Heleioporus psammophilus</i> Lee & Main	+	RE
<i>Heleioporus albopunctatus</i> Gray	n/a	RE ¹
<i>Neobatrachus pelobatooides</i> Werner	+	RE
<i>Crinia georgiana</i> Tschudi	+	RE
<i>Ranidella glauerti</i> Loveridge	+	RE
<i>Ranidella pseudinsignifera</i> Main	+	RE
<i>Ranidella insignifera</i> Moore	n/a	RE ¹
<i>Ranidella subinsignifera</i> Littlejohn	+	LE ²
<i>gen. et sp. nov</i> Roberts <i>et al.</i>	+	LE
<i>Geocrinia rosea</i> Harrison	+	LE
<i>Geocrinia lutea</i> Fletcher	+	LE
<i>Geocrinia alba</i> Wardell-Johnson & Roberts	+	LE
<i>Geocrinia vitellina</i> Wardell-Johnson & Roberts	-	LE
<i>Geocrinia lei</i> Fletcher	+	RE
<i>Metacrinia nichollsi</i> Harrison	+	RE
<i>Myobatrachus gouldii</i> Gray	n/a	RE ¹
<i>Pseudophryne guentheri</i> Boulenger	+	RE

¹ may be found within the boundary of the Warren Bioregion, but only peripherally; reservation status in this bioregion is therefore not applicable (n/a). ² found in the lower south-west, a distribution broadly approximating the Warren bioregion.

vitellina and the sunset frog (Myobatrachidae). All four *Geocrinia* species and the sunset frog have highly limited geographic distributions with *G. vitellina* and *G. alba* formally gazetted in the schedules of the Commonwealth's Endangered Species Protection Act, 1992.

The Reserve System

The Warren Bioregion extends over 1 042 000 ha with approximately 25% of that area held as national parks or nature reserves (A-class). We estimate that these reserves incorporate 86% of the aquatic faunal elements found in the region, but 7% of locally restricted species are apparently not included (Table 4). The further inclusion of locally endemic species or assemblages into the reserve system is made difficult because they are often rare as well as restricted in their distribution. Large-scale surveys across all aquatic habitats within the lower south-west would be necessary to locate all of these aquatic fauna. This is unlikely to occur in the near future and so the ability to predict the occurrence of these fauna at unsurveyed sites would be advantageous.

Predictable patterns of faunal communities often occur in wetlands with similar physical and chemical characteristics (*e.g.* Edward *et al.* 1994). Thus, it may be possible to give priority for reservation to aquatic habitats with particular biophysical traits which, elsewhere, support high levels of endemic species. Highly acidic environments and aquatic habitats associated with granite outcrops provide good examples. The four species of copepod, and two species of cladoceran found by Bayly (1992) were all acidophilic and were collected from ponds with low pH. Two species of ostracod, *Ilyodromus candonites* and *Kapcyridopsis asymmetra*, and one species of chironomid, *Allotrissocladius* sp. which are endemic to the Warren Bioregion, have been found in temporary pools associated with granite outcrops (Bayly 1982). In addition, the sunset frog, which is thought to have originated approximately 30-36 million years ago was recently found in an organic-rich swamp at the base of a granite outcrop near Walpole (Roberts *et al.* 1997).

Granite outcrops and other areas elevated above the level of the Eocene marine incursion, such as headwater regions, permanent freshwater flows, and elevated coastal locations receiving orographic rainfall, often support relictual fauna (Main & Main 1991; Hopper *et al.* 1996). These relictual habitats generally provide organically rich and permanently moist microhabitats (Main & Main 1991) for a fauna which has persisted since eustatic changes and the onset of seasonal aridity in the early Tertiary (Keast 1981). Relictual fauna are of exceptional importance from a nature conservation perspective and thus the systematic evaluation of existing reserves and

Table 4

Summary of the distribution and reservation status of aquatic fauna found in the Warren Bioregion. Number of species found, thus far, in nature reserves (A-class) or national parks within the region are shown in parentheses.

Taxa	Species ¹	Locally Endemic	Regionally Endemic	Widespread
Invertebrates	156	10 (9)	49 (41)	97 (80)
Amphibians	22	6 (5)	13 (13)	0
Fish ²	14	1 (1)	9 (9)	4 (4)

¹ fully described species only; ² native species only.

the inclusion of new reserves to ensure adequate representation of relictual habitats should be an urgent task.

Are the fauna protected in the reserve system?

National parks and nature reserves provide a refuge for a substantial proportion of the aquatic fauna. Their protection, however, cannot be assured unless on-reserve management procedures and activities outside the reserve system are sympathetic to the preservation and maintenance of their habitat.

The practice of conducting fuel reduction burns during late spring, summer and autumn, when the soil is dry, may inadvertently threaten some aquatic fauna either directly or through the loss of soil as habitat. Species such as *L. salamandroides* oversummer in the moist substrate of peatlands and shrublands (Pusey 1990), while other species deposit drought resistant eggs. These organically-rich soils burn readily and hence the fauna within the substratum may be lost. In addition, there may be changes in local hydrology, such as the creation of more surface pools, or improved drainage of sandy soils, which alter the proportional occurrence of habitats and some aquatic species.

Activities occurring within catchments of conservation reserves may impact adversely on the fauna. Harvey (1996) noted that Poarginup Swamp, within the Lake Muir Nature Reserve, was threatened with increased salinization as a result of nearby agricultural clearing. The swamp is the type locality for two species of water mite, *Acercella poarginup* and *Pseudohydryphantes doegi*, both of which were thought to be extinct as a result of recent hydrological changes occurring within the swamp (Harvey 1996). The latter species, however, has been found in one other location, within the Shannon River National Park (Horwitz 1994).

Adverse land-use activities within catchments are even more apparent in flowing waters where anthropogenic activity upstream may directly influence downstream habitats (e.g. Walker 1985; Davey *et al.* 1987; Campbell & Doeg 1989). Land clearing for agriculture in Western Australia has resulted in increased salinity (Schofield 1990) and sedimentation (Williams 1992) of many rivers and clear-fell logging which occurred in the last decade is still affecting streams today (Growth & Davis 1991; Trayler & Davis, *unpubl. obs.*). A river and stream zone system was introduced into the State Forest in the mid-70s (Anon 1977). This was later modified and today this system is estimated to include some 63 100 ha of land in the southern forest region (Anon 1992a). In addition to increasing the size of the conservation estate, these zones of undisturbed vegetation are designed to act as a buffer and minimize the effect of logging operations on the water quality of nearby streams and rivers (Anon 1992a). Large buffer zones (100 m) have been proven effective in reducing the input of sediment into second order streams (Borg *et al.* 1987), but the effectiveness of smaller zones (20-30 m), which are routinely used on these streams, has not been assessed in Western Australia. While there is ample evidence to suggest that these buffers may prevent increased sedimentation (Davies & Nelson 1994; Clinnick 1985), it is unlikely that

such small buffers would prevent increases in stream salinity. Borg *et al.* (1987) found that even 100 m buffers would not prevent a rise in salinity in streams adjacent to logged coupes. There is, however, some evidence to suggest that a large buffer will reduce the period that salinities remain elevated from fifteen years, as estimated by (Borg *et al.* 1988), to eight years, as documented by Growth & Davis (1991). In the past, rising salinity has not been considered important in high rainfall areas where salinities in logged catchments generally remain within the range considered acceptable for drinking water (Anon 1992b). There is, however, increasing evidence that the invertebrate fauna of this region may be intolerant of relatively small increases in salinity (Growth & Davis 1991; Trayler & Davis, *unpubl. obs.*).

The river and stream zone system plays an important role in the conservation of the endemic invertebrate fauna of lotic origin. Many of these fauna have, thus far, not been found elsewhere in the conservation estate (Table 1) and it is therefore essential that riparian buffer zones adjacent to logging coupes are of an adequate size and that these areas are managed properly. Growth (1992) argued that these buffer strips would be compromised if poorly constructed roads or accessways crossed streams to access coupes. In addition, the headwater streams within the karri forest often comprise low gradient, marshy areas with ill-defined and ephemeral water courses which may not be recognised or mapped as first-order streams. There is, therefore, potential for these areas to be overlooked and not included as part of the stream reserve system.

Translocated and exotic fish species are widespread in the waterways of the south-west with increasing anecdotal evidence that these introduced species have a serious impact on the distribution of the native fish fauna (see Morgan *et al.* 1996). Of particular concern is the translocation of piscivorous species such as the golden perch *Macquaria ambigua* and the silver perch *Bidyanus bidyanus* to private dams within the catchment of the D'Entrecasteaux National Park (Morgan *et al.* 1996). These species might further restrict populations of the native fishes if they were to escape into natural waterways.

The non-aestivating native fish species, *N. balstoni*, *Galaxias occidentalis*, *G. munda*, *Edelia vittata* and *Bostockia porosa* are particularly vulnerable to predation by introduced fishes during summer and autumn dry periods when they are forced to retreat to permanent pools and streams (Morgan *et al.* 1996). With the exception of *N. balstoni*, all of these species as well as the lamprey *Geotria australis*, were once found in abundance in the headwaters of Big Brook near Pemberton (Pen *et al.* 1988, 1991), but today their numbers have declined dramatically (Morgan & Gill 1996). In particular, *G. munda*, which was once extremely common to the system (Pen *et al.* 1988, 1991), has now disappeared upstream of the dam and this has been attributed to the recent introduction of the voracious and piscivorous redfin perch *Perca fluviatilis* to Big Brook Dam (Morgan *et al.* 1996).

Big Brook Dam also provides an ideal habitat and potential refuge for a number of other predatory species including *Salmo trutta*, *Gambusia holbrooki* and

Oncorhynchus mykiss (Morgan *et al.* 1996), whose presence may have also adversely affected the native fish fauna in the Big Brook area. While this is yet to be documented for the Warren Bioregion, elsewhere these fishes have been widely implicated in the fragmentation of fish distributions (*e.g.* Tilzey 1976; Lloyd 1990; Hutchinson 1991; Crowl *et al.* 1992).

A variety of other anthropogenic activities and threatening processes operate within nature reserves in the region. Road building activities and the construction of fire breaks, or scrub rolling for fire suppression, increase the likelihood of sediment deposition into wetland systems and enhance the potential for the spread of soil borne fungal pathogens. The effects of these activities have not been fully documented to date.

Recommendations

Conservation reserves within the Warren Bioregion in south-west of Western Australia play an important role in the protection of both the endemic and cosmopolitan aquatic fauna of the entire south-west. The wetlands within these reserves are important refuges for aquatic invertebrates, fishes and amphibians and will become increasingly so as the pressure of urbanisation and agriculture intensifies. Priority for further reservation should be given to aquatic habitats which are known to support a high level of endemic or relictual fauna. It should, however, be acknowledged that the presence of rare fauna is often difficult to detect, and unique faunal assemblages may occur in many of the undisturbed and unsurveyed wetlands of this region. Thus, all relatively undisturbed aquatic habitats in this region should be afforded some protection, at least until they have been properly surveyed.

Since aquatic habitats do not exist in isolation, they cannot be protected within a reserve system without reference to activities occurring within the catchment. Adverse impacts can arise through land clearing, the construction of dams, the introduction of exotic fishes, habitat removal, salinization, sedimentation and eutrophication associated with agriculture, mining, logging and road construction. Careful planning, which is sensitive to the fragility of aquatic habitats, as well as adequate buffers between anthropogenic activities and reserves are essential for the preservation of the fauna in this region.

The effectiveness of a reserve system is also dependent on adequate management within the reserves themselves, where it is essential that due consideration be given to the habitat requirements of the aquatic fauna. This is particularly important for aquatic environments which may be dry for six months or more each year. Seasonal and ephemeral waterbodies must be recognised as comprising an important part of the diversity of aquatic habitats and the absence of water in these environments does not necessarily preclude the presence of aquatic fauna.

Differences between management policy and practice can lead to detrimental effects on aquatic environments (see Davies & Nelson 1994). The environmental awareness of workers carrying out management procedures may be improved through training

specifically in ecological and environmental issues. This important aspect of the management of natural areas currently receives little attention.

Finally, we cannot afford to be complacent with respect to the conservation of the aquatic fauna. Although reserved lands within the Warren Bioregion are extensive, these areas may not be protected in perpetuity. This has been demonstrated by the recent excision of land from the D'Entrecasteaux National Park, near Lake Jasper for the potential purpose of sand mining.

References

- Allen G R 1982 A Field Guide to Inland Fishes of Western Australia. Western Australian Museum, Perth.
- Anon. 1977 General Working Plan 86. Forests Department, Perth.
- Anon. 1992a Management strategies for the south-west forests of Western Australia: A review. Department of Conservation and Land Management, Perth.
- Anon. 1992b Australian water quality guidelines for fresh and marine waters. National Water Quality Management Strategy. Australian and New Zealand Environment and Conservation Council, Canberra.
- Anon. 1994 Australian Threatened Fishes 1994 supplement. Australian Society of Fish Biology Newsletter 24:19-24.
- Anon. 1994 Guidelines for Protected Area Management Categories. IUCN, Gland, Switzerland.
- Balla S A 1994 Wetlands of the Swan Coastal Plain Vol 1: Their Nature and Management. Water Authority of Western Australia & the Department of Environmental Protection, Perth.
- Bayly I A E 1979 Further contributions to a knowledge of the centropagid genera *Boeckella*, *Hemiboeckella* and *Calamoecia* (athalassic calanoid copepods). Australian Journal of Marine and Freshwater Research 30:103-127.
- Bayly I A E 1982 Invertebrate fauna and ecology of temporary pools on granite outcrops in southern Western Australia. Australian Journal of Marine and Freshwater Research 33:599-606.
- Bayly I A E 1992 The micro-crustacea and physico-chemical features of temporary ponds near Northcliffe, Western Australia. Journal of the Royal Society of Western Australia 75:99-106.
- Benzie J A H 1986 *Daphnia occidentalis*, new species (Cladocera: Daphniidae) from Western Australia: new evidence on the evolution of North American *D. ambigua-D. middendorffiana* group. Journal of Crustacean Biology 6:232-245.
- Benzie J A H 1988 The systematics of Australian *Daphnia* (Cladocera: Daphniidae). Species descriptions and keys. Hydrobiologia 166:95-161.
- Borg H, King P D & Loh I C 1987 Stream and ground water response to logging and subsequent regeneration in the southern forest of Western Australia: Interim results from paired catchment studies. Water Authority of Western Australia WH 34, Perth.
- Borg H, Stoneman G L & Ward C G 1988 The effect of logging and regeneration on groundwater, streamflow and stream salinity in the southern forest of Western Australia. Journal of Hydrology 99:253-270.
- Bunn S E & Davies P M 1992 Community structure of the macroinvertebrate fauna and water quality of a saline river system in south-western Australia. Hydrobiologia 248:143-160.
- Bunn S E, Edward D H & Loneragan N R 1986 Spatial and temporal variation in the macroinvertebrate fauna of streams of the northern jarrah forest, Western Australia: community structure. Freshwater Biology 16:67-91.

- Campbell I C & Doeg T J 1989 Impact of timber harvesting and production on streams: A review. *Australian Journal of Marine and Freshwater Research* 40:519-539.
- Christensen P 1982 The distribution of *Lepidogalaxias salamandroides* and other small freshwater fishes in the lower south-west of Western Australia. *Journal of the Royal Society of Western Australia* 65:131-141.
- Christensen P 1992 The karri forest: Its conservation significance and management. Department of Conservation & Land Management, Perth.
- Churchward H M, McArthur W M, Sewell P L & Bartle G A 1988 Landforms and soils of the south-coasts and hinterland, Western Australia: Northcliffe to Manypeaks. CSIRO Divisional Report 88/1, Perth.
- Clinnick P F 1985 Buffer strip management in forest operations: a review. *Australian Forestry* 48:34-45.
- Crowl T A, Townsend A R & McIntosh A R 1992 The impact of introduced brown and rainbow trout on native fish: the case of Australasia. *Reviews in Fish Biology and Fisheries* 2:217-241.
- Davey G W, Doeg T J & Blyth J D 1987 Changes in benthic sediment in the Thomson River, South-Eastern Australia, during construction of the Thomson Dam. *Regulated Rivers* 1:71-84.
- Davis J A, Rosich R S, Bradley J S, Grows J E, Schmidt L G & Cheal F 1993 Wetlands of the Swan Coastal Plain Vol 6: Wetland classification on the basis of water quality and invertebrate community data. Water Authority of Western Australia and the Environmental Protection Authority, Perth.
- Davies P E & Nelson M 1994 Relationships between riparian buffer widths and the effects of logging on stream habitat, invertebrate community composition and fish abundance. *Australian Journal of Marine and Freshwater Research* 45:1289-1305.
- Edward D H D, Gazey P & Davies P M 1994 Invertebrate community structure related to physico-chemical parameters of permanent lakes of the south coast of Western Australia. *Journal of the Royal Society of Western Australia* 77:51-63.
- Grows I O 1992 Macroinvertebrate community structure in the streams of the southern forests of Western Australia: The influence of seasonality, longitudinal gradients and forestry activities. PhD Thesis, Murdoch University, Perth.
- Grows I O & Davis J A 1991 Differences in the macroinvertebrate communities of streams in clearfelled and undisturbed areas in experimental catchments 8 years after harvesting. *Australian Journal of Marine and Freshwater Research* 42:689-706.
- Grows I O & Davis J A 1994 Longitudinal changes in near-bed flows and macroinvertebrate communities in a Western Australian stream. *Journal of the North American Benthological Society* 13:417-438.
- Grows J E, Davis J A, Cheal F, Schmidt R G, Rosich R S & Bradley S J 1992 Multivariate pattern analysis of wetland invertebrate communities and environmental variables in Western Australia. *Australian Journal of Ecology* 17:275-288.
- Halse S A 1989 Wetlands of the Swan Coastal Plain - past to present. In: Swan Coastal Plain Groundwater Management Conference Proceedings (ed G Lowe). Western Australian Water Resources Council, Perth, 105-112.
- Harvey M 1996 A review of the water mite family Pionidae in Australia (Acarina: Hygrobatoidae). *Records of the Western Australian Museum* 17:361-393.
- Hopper S D, Keighery G J & Wardell-Johnson G 1992 Flora of the karri forest and other communities in the Warren Botanical Subdistrict of Western Australia. In: Research on the Impact of Forest Management in South-West Western Australia (ed M Lewis). Department of Conservation and Land Management Occasional Paper 2/92, 1-32.
- Hopper S D, Harvey M, Chappel J, Main A R & Main B Y (1996) Gondwanan heritage - A review. In: The Gondwanan Heritage: Past, Present and Future of the Western Australian Biota. (eds S D Hopper, J Chappel, M Harvey & A George). Surrey Beatty & Sons and Kings Park and Botanic Garden, Chipping Norton and Perth, 1-46.
- Horwitz P 1994 Patterns of endemism in the freshwater fauna of the far southern peatlands and shrublands of south-western Australia. Report. Australian Heritage Commission, Perth.
- Horwitz P 1996 Comparative endemism and richness of the aquatic invertebrate fauna in peatlands and shrublands of far south-western Australia. *Memoirs of the Museum of Victoria* 56:313-322.
- Hutchinson M J 1991 Distribution patterns of redfin perch *Perca fluviatilis* Linnaeus and western pygmy perch *Edelia vittata* Castelnau in the Murray River system Western Australia. *Records of the Western Australian Museum* 15:295-301.
- Jaensch R P 1992 Fishes in wetlands on the south coast of Western Australia. Technical Paper. Department of Conservation and Land Management, Perth.
- Jarvis N T 1979 Western Australia: An atlas of human endeavour. Education and Lands and Surveys Departments of Western Australia, Perth.
- Keast A 1981 Distributional patterns, regional biotas, and adaptations in the Australian biota: A synthesis. In *Ecological Biogeography of Australia* (ed A Keast). Dr W Junk, The Hague.
- Lloyd L 1990 Ecological interactions of *Gambusia holbrooki* with Australian native fishes. In: *Introduced and Translocated Fishes and their Ecological Effects* (ed D.A. Pollard). Australian Society for Fish Biology, Bureau of Rural Resources Proceedings. Australian Government Publishing Service, Canberra, 94-97.
- Main A R 1965 Frogs of Southern Western Australia. Handbook 8, Western Australian Naturalists' Club, Perth.
- Main A R & Main B Y 1991 Report on the Southern Forest Region of Western Australia. Report. Australian Heritage Commission, Perth.
- Morgan D, Gill H & Potter I C 1996 The distribution of freshwater fish in the south-western corner of Western Australia. Report. Water Authority of Western Australia, Perth.
- Morgan D & Gill H 1996 The effect of Big Brook Dam during drought on the fish communities of the Lefroy and Big Brooks. Report. Water and Rivers Commission, Perth.
- Morton D W 1990 Revision of the Australian Cyclopidae (Copepoda: Cyclopoida). II. *Eucyclops* Claus and *Ectocyclops* Brady. *Australian Journal of Marine and Freshwater Research* 41:657-675.
- Mulcahy W M, Churchward H M & Dimmock G M 1972 Landforms and soils on an uplifted peneplain in the Darling Range, Western Australia. *Australian Journal of Soil Research* 10:1-14.
- Olsen G & Skitmore E 1991 State of the Rivers of the South West Drainage Division. Western Australian Water Resources Council 2/91, Perth.
- Pen L J, Hilliard R W & Potter I C 1988 The Effect of Big Brook Dam and the Pemberton Weir on the migration of native fish. Report. Water Authority of Western Australia, Perth.
- Pen L J, Hilliard R W & Power G W 1991 Fish and lamprey monitoring in the Lefroy/Big Brook system in south-west Western Australia. Report. Water Authority of Western Australia, Perth.
- Pusey B J 1990 Seasonality, aestivation and the life history of the salamanderfish *Lepidogalaxias salamandroides* (Pisces: Lepidogalaxiidae). *Environmental Biology of Fishes* 29:15-26.
- Pusey B J & Edward D H 1990a Limnology of the southern acid peat flats, South-Western Australia. *Journal of the Royal Society of Western Australia* 73:29-46.
- Pusey B J & Edward D H 1990b Structure of fish assemblages in waters of the southern acid peat flats, south-western Australia. *Australian Journal of Marine and Freshwater Research* 41: 721-734.
- Roberts J D, Horwitz P, Wardell-Johnson G, Maxson L R & Mahony M 1997 Taxonomy, relationships and conservation of

- a new genus and species of myobatrachid frog from the high rainfall region of south-western Australia. *Copeia* 1997: 373-381.
- Schofield N J 1990 Water Interactions with land use and climate in south western Australia. Western Australian Water Authority Report WS60, Perth.
- Storey A W, Bunn S E, Davies P M & Edward D H 1990 Classification of the macroinvertebrate fauna of the river systems in southwestern Australia in relation to physical and chemical parameters. *Regulated Rivers: Research & Management* 5:217-232.
- Thackway R & Cresswell I D 1995 An interim biogeographic regionalisation for Australia: A framework for setting priorities in the national reserves system cooperative program. Australian Nature Conservation Agency, Canberra.
- Tilzey R D 1976 Observations of interactions between indigenous Galaxiidae and introduced Salmoniidae in the Lake Eucumbene catchment, New South Wales. *Australian Journal of Marine and Freshwater Research* 27:551-564.
- Tyler M J, Smith L A & Johnstone R E 1994. *Frogs of Western Australia*. Western Australian Museum, Perth.
- Walker K F 1985 A review of the ecological effects of river regulation in Australia. *Hydrobiologia* 125:111-129.
- Williams P 1992 The state of Western Australian rivers. *Land and Water Research News* 13:8-15.
- Williams W D, Taaffe R G & Boulton A J 1991 Longitudinal distribution of macroinvertebrates in two rivers subject to salinization. *Hydrobiologia* 210: 151-161.